

Tipranavir disodium



Drug Description

Tipranavir (TPV) is a sulfonamide-containing dihydropyrene. [1]

HIV/AIDS-Related Uses

TPV is a nonpeptidic protease inhibitor (NPI) that has a uniquely robust resistance profile, demonstrating viral load responses against multiple protease inhibitor (PI) resistant HIV-1 strains both in vitro and in clinical studies.[2]

Ritonavir-boosted tipranavir (TPV/r) is currently being evaluated in treatment-experienced adult patients in Phase III clinical trials.[3] Pediatric formulations are currently being evaluated in HIV-infected patients age 2 to 18 years in Phase I, II, and III clinical trials. No results have been reported to date.[4] [5]

Pharmacology

TPV flexibly binds to the active site of HIV-1 protease, suppressing viral replication.[6] [7]

In a Phase II trial of TPV in PI-experienced patients, TPV/r (500/200 mg dose) consistently produced TPV trough plasma levels in excess of 10 times the IC₉₀ (the concentration of drug required to inhibit viral replication by 90%) for PI-resistant HIV-1. Steady state plasma concentrations were reached within the first 7 days of treatment with twice-daily dosing.[8]

TPV/r demonstrated a greater than 10-fold reduction in viral load in treatment-experienced patients with multiple PI-resistant HIV.[9] In treatment-naïve patients, TPV/r regimens demonstrated a 30-fold HIV RNA reduction after 2 weeks of treatment.[10]

A minimum of 16 to 20 protease gene mutations may be required for reduced susceptibility to TPV.[11] [12] Reduced susceptibility was demonstrated only in isolates with three or more universal protease-associated mutations (UPAMs) at codons L33I/V/F, V82A/F/L/T, I84V, and/or L90M, in addition to other protease gene mutations. Few isolates with only one or two UPAMs

displayed reduced susceptibility to TPV; however, one or two of these mutations caused reduced susceptibility to available PIs. A breakpoint for TPV susceptibility was observed at an IC₅₀ of approximately twofold wild type.[13]

TPV is a substrate and inducer of cytochrome P450 CYP3A in animals and humans. TPV/r acts as a CYP3A4 inhibitor due to the potent inhibitory effect of ritonavir.[14] In rats, glucuronidation of TPV is the major clearance mechanism.[15]

Adverse Events/Toxicity

The most common adverse events seen in TPV doses up to 1200 mg were mild to moderate nausea, vomiting, and diarrhea or loose stools. Adverse events were similar among all dosage regimens.[16] In a Phase II clinical trial of 216 patients, 15.3% experienced diarrhea of Grade 2 or greater, and 11.6% experienced vomiting; however, no clear dose response relationship was observed. A dose-dependent trend was observed for Grade 3 and 4 adverse events, laboratory abnormalities, and adverse event-related treatment discontinuations. Eleven patients (5.1%) experienced serious adverse events during the first 4 weeks of therapy; two were regarded as treatment-related.[17] In a phase II clinical trial of 296 patients on TPV/r alone or in combination with another PI, the frequency of adverse effects and laboratory abnormalities were similar between all arms after 4 weeks of therapy. Triglyceride elevations were the most frequent laboratory abnormality.[18]

Drug and Food Interactions

TPV decreases the steady state plasma concentrations of didanosine (ddI), stavudine, lamivudine, and zidovudine. However, the observed differences (15% to 46% decrease) in nucleoside reverse transcriptase inhibitor (NRTI) concentrations are not of clinical importance, and TPV can be combined with these agents.[19] Coadministration of TPV with enteric-coated (EC) ddI results in a 32% increase in TPV peak plasma concentration (C_{max}) and a 34% decrease in TPV plasma trough concentration (C_{12h}). EC ddI should be administered 4 hours apart from TPV/r.[20]

Tipranavir disodium

Drug and Food Interactions (cont.)

Antacids reduce the absorption of TPV by 25% to 29%, requiring adjustment to the timing of antacid use. TPV increases atorvastatin plasma concentrations over ninefold, suggesting clinicians should carefully monitor patients on this combination for atorvastatin-associated adverse effects.[21]

Clinical Trials

For information on clinical trials that involve Tipranavir disodium, visit the ClinicalTrials.gov web site at <http://www.clinicaltrials.gov>. In the Search box, enter: Tipranavir disodium AND HIV Infections.

Dosing Information

Mode of Delivery: Oral.[22]

Dosage Form: Phase II clinical trials have evaluated TPV in a 300 mg hard-gel capsule. This formulation, which required high doses and a high pill burden to obtain optimal plasma levels, was replaced with a 250 mg soft-gel capsule available in a Self-Emulsifying Drug Delivery System.[23] [24] Phase II clinical trials have evaluated TPV/r at 500/100 mg, 500/200 mg, and 750/200 mg twice daily doses. Of these, the 500/200 mg dose has been selected for further study in Phase III trials based on optimal pharmacokinetic, safety, and tolerability data.[25] [26]

Storage: Store capsules at 2 C to 8 C in a cool, dry place protected from light, heat, and moisture. TPV may become less active in such conditions, and the soft-gel capsules may stick together in warm, moist conditions.[27]

Chemistry

CAS Name: 2-Pyridinesulfonamide, N-(3-[(1R)-1-[(6R)-5,6-dihydro-4-hydroxy-2-oxo-6-(2-phenylethyl)-6-propyl-2H-pyran-3-yl]propyl]phenyl)-5-(trifluoromethyl)-, disodium salt[28]

CAS Number: 191150-83-1[29]

Molecular formula: C₃₁H₃₁F₃N₂Na₂O₅S[30]

C 57.6%, H 4.8%, F 8.8%, N 4.3%, Na 7.1%, O 12.4%, S 5.0%[31]

Molecular weight: 646.64[32]

Other Names

PNU 140690E[33]

TPV[34]

PNU 140690[35]

Further Reading

[No authors listed] Tipranavir reduces viral load. AIDS Patient Care STDS. 2002 Sep;16(9):462.

Larder BA, Hertogs K, Bloor S, van den Eynde CH, DeCian W, Wang Y, Freimuth WW, Tarpley G. Tipranavir inhibits broadly protease inhibitor-resistant HIV-1 clinical samples. AIDS. 2000 Sep 8;14(13):1943-8. PMID: 10997398

McCallister S, Valdex H, Curry K, MacGregor T, Borin M, Freimuth W, Wang Y, Mayers DL. A 14-Day Dose-Response Study of the Efficacy, Safety, and Pharmacokinetics of the Nonpeptidic Protease Inhibitor Tipranavir in Treatment-Naive HIV-1-Infected Patients. J Acquir Immune Defic Syndr. 2004 Apr 20;35(4):376-82. PMID: 15097154

Schake D. How flexible is tipranavir in complex with the HIV-1 protease active site? AIDS. 2004 Feb 20;18(3):579-80. PMID: 15090819

Yeni P. Tipranavir: a protease inhibitor from a new class with distinct antiviral activity. J Acquir Immune Defic Syndr. 2003 Sep;34 Suppl 1:S91-4. PMID: 14562864

Tipranavir disodium

Manufacturer Information

Tipranavir disodium
Boehringer Ingelheim GmbH
55216 Ingelheim am Rhein
Germany

For More Information

Contact your doctor or an AIDSinfo Health
Information Specialist:

- Via Phone: 1-800-448-0440 Monday - Friday,
12:00 p.m. (Noon) - 5:00 p.m. ET
- Via Live Help: http://aidsinfo.nih.gov/live_help
Monday - Friday, 12:00 p.m. (Noon) - 4:00 p.m. ET

References

1. Antimicrob Agents Chemother - 1997 May;41(5):1058-63
2. Conf Retroviruses Opportunistic Infect. - 10th, 2003. Abstract 179.
3. Conf Retroviruses Opportunistic Infect. - 10th, 2003. Abstract 596.
4. ClinicalTrials.gov - Tipranavir in Combination with Ritonavir in HIV-Infected Children. Available at: <http://www.clinicaltrials.gov/ct/show/NCT00076999>. Accessed 09/07/04.
5. ClinicalTrials.gov - Tipranavir/Ritonavir in HIV Patients with Limited Treatment Options. Available at: <http://www.clinicaltrials.gov/ct/show/NCT00062660>. Accessed 09/07/04.
6. Antimicrob Agents Chemother - 1997 May;41(5):1058-63
7. AIDS - 2000;14(3):1943-1948
8. Conf Retroviruses Opportunistic Infect. - 10th, 2003. Abstract 528.
9. Conf Retroviruses Opportunistic Infect. - 10th, 2003. Abstract 596.
10. Conf Retroviruses Opportunistic Infect. - 7th, 2000. Abstract 673.
11. Intl Wkshp Manage Treatment-Experienced Patients - 2nd, 2002. Abstract P5.
12. Intl HIV Drug Resistance Wkshp - 11th, 2002. Abstract 127.
13. Conf Retroviruses Opportunistic Infect. - 10th, 2003. Abstract 596.
14. IAS Conf on HIV Pathogenesis and Treatment - 2nd, 2003. Abstract 865.
15. Conf Retroviruses Opportunistic Infect. - 7th, 2000. Abstract 81.
16. Conf Retroviruses Opportunistic Infect. - 7th, 2000. Abstract 673.

Tipranavir disodium



17. Conf Retroviruses Opportunistic Infect. - 10th, 2003. Abstract 179.
18. Intl AIDS Conf - 15th, 2004. Abstract WeOrB1236.
19. Conf Retroviruses Opportunistic Infect. - 7th, 2000. Abstract 81.
20. IAS Conf on HIV Pathogenesis and Treatment - 2nd, 2003. Abstract 865.
21. Intl Wkshp on Clinical Pharmacology of HIV Therapy - 5th, 2004. Abstract 5.2.
22. Conf Retroviruses Opportunistic Infect. - 10th, 2003. Abstract 528.
23. Conf Retroviruses Opportunistic Infect. - 10th, 2003. Abstract 528.
24. European AIDS Conf - 9th, 2003. Abstract 7.2/3.
25. Conf Retroviruses Opportunistic Infect. - 10th, 2003. Abstract 528.
26. IAS Conf on HIV Pathogenesis and Treatment - 2nd, 2003. Abstract 865.
27. Ontario HIV Pharmacy Professional Specialty Group - 2003. Tipranavir Fact Sheet. Available at: <http://www.tthhivclinic.com>. Accessed 9/27/04.
28. ChemIDplus - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/chemidlite.jsp>. Accessed 10/1/04.
29. USPD - 2003; p. 861
30. USPD - 2003; p. 861
31. Calculation. -
32. USPD - 2003; p. 861
33. USPD - 2003; p. 861
34. NIAID HIV/OI Therapeutics Database - http://apps1.niaid.nih.gov/struct_search/an/an_search.htm. Accessed 9/8/04.
35. ChemIDplus - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/chemidlite.jsp>. Accessed 9/30/04.